Root **ICROSENSE**



Made Germany



ROOT ENHANCEMENT

Germination, rooting and the adaptation to the field can be streamlined with MICROSENSE® Root. Microbial organisms producing natural hormones such as Auxin and Gibberellin are combined with natural organic acids, amino acids and fulvic and humic acids to accelerate seed germination, boost root length and density and raise rhizospheric activity. Amino acids such as Lysine, Valine, Serine, Tryptophan and Glutamate optimize rooting in plants produced from seeds and cuttings. Improving germination and rooting affects nutrient uptake efficiency, accelerates growth rate and leads to higher yields.

APPLICA BENEFITS & RECOMMENDED 0

0.15 - 0.5 L / Ha via spaying

0.30 - 0.50 L / m³

1 L / 1 ton seed

- Optimizes seed germination and root development from cuttings
- Increases vigor of seed germination
- Reduces soil compaction causing germination difficulties
- Minimizes damages caused by low temperature
- Leads higher yields
- Accelerates plant growth

- Increases fertilizer use efficiency with less nutrient leaching
 - Promotes root respiration and vertical development
- Improves soil quality
- Regenerates degraded soil
- Increases soil biodiversity and biological activity in the rhizosphere
- Boosts root length and density

Total amount/vegetation period 4 – 6 L / Ha (1 – 2 L / Ha per application)

FIELDS OF APPLICATION **RECOMMENDED APPLICATION RATES***

Soil

Foliar

Seeds

Substrates

Hydroponics

- Open field and Greenhouse
- Agriculture
- Vegetable Production
- Fruit Production
- Substrate Cultivation
- Turf and Landscaping
- Seed Treatment

* Standard recommendations that can vary according to soil properties, cultivated crop and local application conditions.

20 - 40 ml / 1.000 L nutrient solution during cultivation cycle

SHAKE WELL BEFORE USE

Please remind to shake well the bottle or container prepared mix well before use.



Before use in combination with other agents, first preparation of a sample mix in a small scale is recommended to avoid any risk of flocculation.

BLEND USE NOTICE

TÜΛ

PREPARATION OF MIXTURE

Make sure sufficient amount of solution is added to half water filled tank while agitator is running. Add additional substances or crop protection agents before filling the tank fully with water.



ן י		L	
と	ń	L	
)	

INHERENT COMPOSITION* (Typical values based on Fresh Matter)

Humic Acid	10 %	Nitrogen (N)	2 %
Total Amino Acid	7 %	Zinc (Zn)	1.5 %
Azosiprillum Brasselience	1x10 ⁷	рН	4 – 5
Azotobacter Chrococcum	1x10 ⁷	Density	1.15 – 1.16 g/cm ³

German technology

* Humintech keeps the right to change the composition of the product.



Administration & Distribution: HUMINTECH GMBH Am Pösenberg 9 – 13 • 41517 Grevenbroich / Germany • E-Mail: info@humintech.com • Internet: www.humintech.com



Store in a dry area, protected from frost, heat and direct sunlight. Can be stored up to 2 years in its original packaging. **KEEP CONTAINER CLOSED DURING STORAGE!**

KEEP OUT OF THE REACH OF CHILDREN, PETS AND FOOD



